

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**


**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

### Amendment to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application.

What is claimed is:

1. (currently amended): A compound of Formula I



Formula I

wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  are the same or different and each is hydrogen, halogen, alkyl of 1 to 24 carbon atoms, haloalkyl of 1 to 24 carbon atoms, aryl of 6 to 24 carbon atoms, alkenyl of 2 to 24 carbon atoms, acyl of 1 to 24 carbon atoms, aralkyl of 7 to 24 carbon atoms, aracyl of 6 to 24 carbon atoms; OR,  $NRR'$ ,  $CONRR'$ ,  $OCOR$ , CN, SR,  $SO_2R$ ,  $SO_3H$ ,  $SO_3M$ , wherein M is an alkali metal, R and  $R'$  are the same or different and each is hydrogen, alkyl of 1 to 24 carbon atoms, haloalkyl of 1 to 24 carbon atoms, aryl of 6 to 24 carbon atoms, alkenyl of 2 to 24 carbon atoms, acyl of 1 to 24 carbon atoms, cycloalkyl of 1 to 24 carbon atoms, cycloacyl of 5 to 24 carbon atoms, aralkyl of 7 to 24 carbon atoms, or aracyl of 6 to 24 carbons atoms, and optionally with either of  $R_1$  and  $R_2$ ,  $R_3$  and  $R_4$ , or  $R_4$  and  $R_5$ , taken together being a part of a saturated or unsaturated fused carbocyclic ring optionally containing O, N, or S atoms in the ring;

each of T and T' is independently a direct bond, oxygen, NR, sulfur or a functional groups containing these elements;

X is independently selected from hydrogen and a blocking group selected from one or more of the following groups: allyl, —COR<sup>a</sup>, —SO<sub>2</sub>R<sup>b</sup>, —SiR<sup>c</sup>R<sup>d</sup>R<sup>e</sup>, —PR<sup>f</sup>R<sup>g</sup>, —POR<sup>f</sup>R<sup>g</sup> and —CONHR<sup>h</sup>,

wherein

each R<sup>a</sup> is independently selected from C<sub>1</sub>—C<sub>8</sub> alkyl, halogen-substituted C<sub>1</sub>—C<sub>8</sub> alkyl, C<sub>5</sub>—C<sub>12</sub> cycloalkyl, C<sub>2</sub>—C<sub>8</sub> alkenyl, —CH<sub>2</sub>—CO—CH<sub>3</sub>, C<sub>1</sub>—C<sub>12</sub> alkoxy, and phenyl or phenoxy which is unsubstituted or substituted by C<sub>1</sub>—C<sub>12</sub> alkyl, C<sub>1</sub>—C<sub>4</sub> alkoxy, halogen and/or benzyl;

each R<sup>b</sup> is independently selected from C<sub>1</sub>—C<sub>12</sub> alkyl, C<sub>6</sub>—C<sub>10</sub> aryl and C<sub>7</sub>—C<sub>18</sub> alkylaryl;


each R<sup>c</sup>, R<sup>d</sup> and R<sup>e</sup> is independently selected from C<sub>1</sub>—C<sub>18</sub> alkyl, cyclohexyl, phenyl and C<sub>1</sub>—C<sub>18</sub> alkoxy;


each R<sup>f</sup> and R<sup>g</sup> is independently selected from C<sub>1</sub>—C<sub>12</sub> alkoxy, C<sub>1</sub>—C<sub>12</sub> alkyl, C<sub>5</sub>—C<sub>12</sub> cycloalkyl, and phenyl or phenoxy which is unsubstituted or substituted by C<sub>1</sub>—C<sub>12</sub> alkyl, C<sub>1</sub>—C<sub>4</sub> alkoxy, halogen and/or benzyl; and

each R<sup>h</sup> is independently selected from C<sub>1</sub>—C<sub>8</sub> alkyl, C<sub>5</sub>—C<sub>12</sub> cycloalkyl, C<sub>2</sub>—C<sub>8</sub> alkenyl, —CH<sub>2</sub>—CO—CH<sub>3</sub>, and phenyl which is unsubstituted or substituted by C<sub>1</sub>—C<sub>12</sub> alkyl, C<sub>2</sub>—C<sub>8</sub> alkenyl, C<sub>1</sub>—C<sub>4</sub> alkoxy, halogen and/or benzyl;


each of Y, R<sub>6</sub> and R<sub>7</sub> are each independently hydrogen, hydrocarbyl group, a functional hydrocarbyl group, halogen, hydroxyl, cyano, -O(hydrocarbyl), -O(functional hydrocarbyl), -N(hydrocarbyl)<sub>2</sub>, -N(functional hydrocarbyl)<sub>2</sub>, -N(hydrocarbyl)(functional hydrocarbyl), -S(hydrocarbyl), -S(functional hydrocarbyl), -SO<sub>2</sub>(hydrocarbyl), -SO<sub>2</sub>(functional hydrocarbyl), -SO<sub>3</sub>(hydrocarbyl), -SO<sub>3</sub>(functional hydrocarbyl), -CO<sub>2</sub>(hydrocarbyl), -CO<sub>2</sub>(functional hydrocarbyl), -CO(hydrocarbyl), -CO(functional hydrocarbyl), -OCO(hydrocarbyl), -OCO(functional hydrocarbyl), -CONH<sub>2</sub>, -CONH(hydrocarbyl), -CONH(functional hydrocarbyl), -CON(hydrocarbyl)<sub>2</sub>, -CON(hydrocarbyl)(functional hydrocarbyl), -CON(functional hydrocarbyl)<sub>2</sub>, wherein the hydrocarbyl or functional hydrocarbyl may be the same or different and has 1 to 24 carbon atoms;

Z is Y,





or




and wherein L is ~~hydrogen, a hydrocarbyl of 1 to 24 carbon atoms, or a functional hydrocarbyl of 1 to 24 atoms;~~ selected from the group consisting of: hydrogen; an alkyl of 1 to 24 carbon atoms optionally substituted by one or more hydroxy, alkoxy, carboxy, carboalkoxy, amino, amido, carbamato, or epoxy groups, and which may contain one or more carbonyl groups, oxygen atoms or nitrogen atoms in the chain; an alkenyl of 2 to 24 carbon atoms optionally substituted by one or more hydroxy, alkoxy, carboxy, carboalkoxy, amino, amido, carbamato, or epoxy groups, and which may contain one or more carbonyl groups, oxygen atoms or nitrogen atoms in the chain; an aralkyl of 7 to 24 carbon atoms optionally substituted by one or more hydroxy, alkoxy, chloro, cyano, carboxy, carboalkoxy, amino, amido, carbamato, or epoxy groups, and which may contain one or more carbonyl groups, oxygen atoms or nitrogen atoms in the chain.

2. (currently amended): A compound according to claim 1 wherein T' is an oxygen atom, ~~and Y is L.~~

3. (cancelled).


4. (currently amended): A compound according to claim 41 wherein TZ is




~~and wherein L is hydrogen, a hydrocarbyl group of 1 to 24 carbon atoms, or a functional hydrocarbyl group of 1 to 24 carbon atoms;~~  
~~X is independently selected from hydrogen and a blocking group; and~~

$R_6$  and  $R_7$  are independently hydrogen, hydrocarbyl, functional hydrocarbyl, halogen, hydroxyl,  $-\text{O}(\text{hydrocarbyl})$ ,  $-\text{O}(\text{functional hydrocarbyl})$ ,  $-\text{S}(\text{hydrocarbyl})$ ,  $-\text{SO}_2(\text{hydrocarbyl})$ ,  $-\text{SO}_3(\text{hydrocarbyl})$ ,  $-\text{COO}(\text{hydrocarbyl})$ ,  $-\text{CO}(\text{hydrocarbyl})$ ,  $-\text{OCO}(\text{hydrocarbyl})$ ,  $-\text{N}(\text{hydrocarbyl})(\text{hydrocarbyl})$ ,  $-\text{S}(\text{functional hydrocarbyl})$ ,  $-\text{SO}_2(\text{functional hydrocarbyl})$ ,  $-\text{SO}_3(\text{functional hydrocarbyl})$ ,  $-\text{COO}(\text{functional hydrocarbyl})$ ,  $-\text{CO}(\text{functional hydrocarbyl})$ ,  $-\text{OCO}(\text{functional hydrocarbyl})$ ,  $-\text{N}(\text{functional hydrocarbyl})(\text{functional hydrocarbyl})$  or cyano.

5. (original): A compound according to claim 2 wherein  $R_1$  to  $R_7$  and X are hydrogen, and TZ is




6. (original): A compound according to claim 1 wherein TZ is:



7. (currently amended): A compound according to claim 6 wherein T' is oxygen, ~~and Y~~  
is ~~L~~

8. (original): A compound according to claim 7 wherein  $R_1$  to  $R_7$  and X are hydrogen.

9. (currently amended): A compound of Formula VI:



(Formula VI)

wherein T, Z, R<sub>1</sub> to R<sub>7</sub>, X are as defined in claim 1, and r is an integer between 2 and 4;

when r is 2, D is selected from the group consisting of C<sub>2</sub>—C<sub>16</sub> alkyl, C<sub>4</sub>—C<sub>12</sub> alkenyl, xylylene, C<sub>3</sub>—C<sub>20</sub> alkyl which is interrupted by one or more oxygen atoms, hydroxy-substituted C<sub>3</sub>—C<sub>20</sub> alkyl which is interrupted by one or more oxygen atoms, —CH<sub>2</sub>CH(OH)CH<sub>2</sub>O—R<sup>15</sup>—OCH<sub>2</sub>CH(OH)CH<sub>2</sub>—, —CO—R<sup>16</sup>—CO—, —CO—NH—R<sup>17</sup>—NH—CO—, —(CH<sub>2</sub>)<sub>s</sub>—COO—R<sup>18</sup>—OCO—(CH<sub>2</sub>)<sub>s</sub>—

a polyoxyalkylene bridge member of the Formula XX

—CH<sub>2</sub>—CH(OH)—CH<sub>2</sub>—O—(CH<sub>2</sub>—(CH<sub>2</sub>)<sub>u</sub>—O—)<sub>mm</sub>—CH<sub>2</sub>—CH(OH)—CH<sub>2</sub>— (XX),


a polyoxyalkylene bridge member of the Formula XXI

—CO—(CH<sub>2</sub>)<sub>u</sub>—O—(CH<sub>2</sub>—(CH<sub>2</sub>)<sub>u</sub>—O—)<sub>mm</sub>—(CH<sub>2</sub>)<sub>u</sub>—CO— (XXI),

a polyoxyalkylene bridge member of the Formula XXII

—YY—O—CO(CH<sub>2</sub>)<sub>u</sub>—O—(CH<sub>2</sub>—(CH<sub>2</sub>)<sub>u</sub>—O—)<sub>mm</sub>—(CH<sub>2</sub>)<sub>u</sub>—COO—YY— (XXII),


a polyoxyalkylene bridge member of the Formula XXIII



a polyoxyalkylene bridge member of the Formula XXIV




a polyoxyalkylene bridge member of the Formula XXV



a polyoxyalkylene bridge member of the Formula XXVI



and a polyoxyalkylene bridge member of the Formula XXVII



wherein  $a + c = 2.5$  and  $b = 8.5$  to  $40.5$  or  $a + c = 2$  to  $33$  and  $b = 0$ ,

$R^{21}$  is hydrogen or  $C_1\text{---}C_{16}$  alkyl,

~~$R^{22}$  is halogen or  $\text{---O---}R^{23}$ ;~~

~~$R^{23}$  is hydrogen,  $C_4\text{---}C_6$  alkyl,  $C_3\text{---}C_6$  alkenyl, aryl, or aryl  $\text{---}C_4\text{---}C_4\text{---}$  alkyl;~~

~~$R^{24}$  is hydrogen,  $C_4\text{---}C_{42}$  alkyl or aryl;~~

~~$R^{25}$  is  $C_4\text{---}C_{46}$  alkyl,  $C_5\text{---}C_{42}$  cycloalkyl,  $C_3\text{---}C_6$  alkenyl,~~

~~$C_4\text{---}C_{42}$  alkylaryl or aryl  $\text{---}C_4\text{---}C_4\text{---}$  alkyl;~~

~~$R^{26}$  is hydrogen or  $C_4\text{---}C_4$  alkyl;~~

~~$R^{27}$  is hydrogen,  $C_4\text{---}C_{48}$  alkyl,  $C_3\text{---}C_6$  alkenyl,  $C_4\text{---}C_{48}$  alkoxy, halogen or aryl  $\text{---}C_4\text{---}$   $C_4\text{---}$  alkyl;~~

~~$R^{28}$  and  $R^{29}$  independently of one another are hydrogen,  $C_4\text{---}C_{48}$  alkyl,  $C_3\text{---}C_6$  alkenyl, or~~

~~$C_4\text{---}C_{48}$  alkoxy, or halogen;~~

~~$R^{30}$  is hydrogen,  $C_4\text{---}C_4$  alkyl or CN;~~

YY is unsubstituted or substituted  $C_2\text{---}C_{20}$  alkyl,

kk is zero or an integer from 1-16,

$B_1$  is O or NH,

mm is an integer from 2 to 60,

nn is an integer from 2 to 6,

u is an integer from 1 to 4;

when r is 3, D is  $\text{---}[(\text{CH}_2)_s\text{---COO-}]_3\text{---}R^{19}$

and when r is 4, D is  $\text{---}[(\text{CH}_2)_s\text{---COO-}]_4\text{---}R^{20}$

wherein  $R^{19}$  is  $C_3\text{---}C_{10}$  alkanetriyl and  $R^{20}$  is  $C_4\text{---}C_{10}$  alkanetetryl; and

s is 1-6;


$R^{15}$  is  $C_2-C_{10}$  alkyl,  $C_2-C_{10}$  oxaalkyl or  $C_2-C_{10}$  dithiaalkyl, phenyl, naphthyl, diphenyl, or  $C_2-C_6$  alkenyl, or phenylene-XX-phenylene wherein XX is  $-O-$ ,  $-S-$ ,  $-SO_2-$ ,  $-CH_2-$ , or  $-C(CH_3)_2-$ ;

$R^{16}$  is  $C_2-C_{10}$  alkyl,  $C_2-C_{10}$  oxaalkyl or  $C_2-C_{10}$  dithiaalkyl, phenyl, naphthyl, diphenyl, or  $C_2-C_6$  alkenyl provided that when r is 3 the alkenyl has at least 3 carbons;

$R^{17}$  is  $C_2-C_{10}$  alkyl, phenyl, naphthyl, diphenyl, or  $C_2-C_6$  alkenyl, methylenediphenylene, or  $C_4-C_{15}$  alkylphenyl; and

$R^{18}$  is  $C_2-C_{10}$  alkyl, or  $C_4-C_{20}$  alkyl interrupted by one or more oxygen atoms.

10. (currently amended): A compound of Formula (VII):



(Formula VII)

wherein T, T', Y, Z,  $R_1$  to  $R_7$ , are as defined in claim 1; r is 2 or 3;

when r is 2,  $X'$  is  $-CO-R^{16}-CO-$ ,  $-CO_2-R^{16}-CO_2-$ ,  $-SO_2-R^{16}-SO_2-$ ,  $-CO-NH-R^{17}-NH-CO-$ , a polyoxyalkylene bridge member of Formula  $-CO-(CH_2)_u-O-(CH_2-(CH_2)_u-O)_{mm}-(CH_2)_u-CO-$ , or  $-COC(R^{21})HCH_2NH(C_{nn}H_{2nn}O)_mC_{nn}H_{2nn}-NHCH_2-C(R^{21})HCO-$

when r = 3,  $X'$  is:



$-(\text{---CO}_2\text{---R}^{16})\text{R}^{19}$ ,  $-(\text{---CONH---R}^{16})\text{R}^{19}$ ,  $-(\text{---SO}_2\text{---R}^{16})\text{R}^{19}$

wherein  $\text{R}^{19}$  is  $\text{C}_3\text{---C}_{10}$  alkanetriyl and

$\text{R}^{16}$  is  $\text{C}_2\text{---C}_{10}$  alkyl,  $\text{C}_2\text{---C}_{10}$  oxaalkyl or  $\text{C}_2\text{---C}_{10}$  dithiaalkyl, phenyl, naphthyl, diphenyl, or  $\text{C}_2\text{---C}_6$  alkenyl provided that when  $r$  is 3 the alkenyl has at least 3 carbons;

$\text{R}^{17}$  is  $\text{C}_2\text{---C}_{10}$  alkyl, phenyl, naphthyl, diphenyl, or  $\text{C}_2\text{---C}_6$  alkenyl, methylenediphenylene, or  $\text{C}_4\text{---C}_{15}$  alkylphenyl; and

~~$\text{R}^{18}$  is  $\text{C}_2\text{---C}_{40}$  alkyl, or  $\text{C}_4\text{---C}_{20}$  alkyl interrupted by one or more oxygen atoms; and~~


$\text{R}^{21}$  is hydrogen or  $\text{C}_1\text{---C}_{16}$  alkyl; and

$mm$  is an integer from 2 to 60,

$nn$  is an integer from 2 to 6,

$u$  is an integer from 1 to 4.

11. (original): A compound of Formula (VIII):




(Formula VIII)

wherein T, T', Y, Z,  $\text{R}_1$  to  $\text{R}_5$ ,  $\text{R}_7$  and X, are as defined in claim 1;


$\text{R}_6$  is selected from the group consisting of straight chain alkyl of 1 to 12 carbon atoms, branched chain alkyl of 1 to 12 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, alkyl substituted by cyclohexyl, alkyl interrupted by cyclohexyl, alkyl

substituted by phenylene, alkyl interrupted by phenylene, benzylidene, —S—, —S—S—, —S—E—S—, —SO—, —SO<sub>2</sub>—, —SO—E—SO—, —SO<sub>2</sub>—E—SO<sub>2</sub>—, —CH<sub>2</sub>—NH—E—NH—CH<sub>2</sub>—, and



wherein E is selected from the group consisting of alkyl of 2 to 12 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, alkyl interrupted by cyclohexyl of 8 to 12 carbon atoms, alkyl terminated by cyclohexyl of 8 to 12 carbon atoms; and r is an integer between 2 and 4

12. (original): A compound of Formula IX



(Formula IX)

wherein T, T' Y, X, and R<sub>1</sub> to R<sub>7</sub> are defined as in claim 1;

r is an integer between 2 and 4;

when r is 2, D is selected from the group consisting of C<sub>2</sub>—C<sub>16</sub>alkylene, C<sub>4</sub>—C<sub>12</sub>alkenylene, xylylene, C<sub>3</sub>—C<sub>20</sub>alkylene which is interrupted by one or more oxygen atoms, hydroxy-substituted C<sub>3</sub>—C<sub>20</sub>alkylene which is interrupted by one or more

oxygen atoms,  $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{O}-\text{R}^{15}-\text{OCH}_2\text{CH}(\text{OH})\text{CH}_2-$ ,  $-\text{CO}-\text{R}^{16}-\text{CO}-$ ,  
 $-\text{CO}-\text{NH}-\text{R}^{17}-\text{NH}-\text{CO}-$ , and  $-(\text{CH}_2)_s-\text{COO}-\text{R}^{18}-\text{OCO}-(\text{CH}_2)_s-$ ; and

when r is 3, D is  $-\text{[-}(\text{CH}_2)_s-\text{COO-}]_3-\text{R}^{19}$

and when r is 4, D is  $-\text{[-}(\text{CH}_2)_s-\text{COO-}]_4-\text{R}^{20}$

wherein  $\text{R}^{19}$  is  $\text{C}_3-\text{C}_{10}$ alkanetriyl and  $\text{R}^{20}$  is  $\text{C}_4-\text{C}_{10}$ alkanetetryl;

s is 1-6;

$\text{R}^{15}$  is  $\text{C}_2-\text{C}_{10}$  alkylene phenylene or a phenylene-x-phenylene- group, wherein X is  
 $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}_2-$ ,  $-\text{CH}_2-$ , or  $-\text{C}(\text{CH}_3)_2-$ ;

$\text{R}^{16}$  is  $\text{C}_2-\text{C}_{10}$  alkylene,  $\text{C}_2-\text{C}_{10}$  oxaalkylene or  $\text{C}_2-\text{C}_{10}$  dithiaalkylene, phenylene,  
naphthylene, diphenylene or  $\text{C}_2-\text{C}_6$  alkenylene;

$\text{R}^{17}$  is  $\text{C}_2-\text{C}_{10}$  alkylene, phenylene, naphthylene, methylenediphenylene or  $\text{C}_7-\text{C}_{15}$   
alkylphenylene, and

$\text{R}^{18}$  is  $\text{C}_2-\text{C}_{10}$  alkylene or  $\text{C}_4-\text{C}_{20}$  alkylene which is interrupted by one or more  
oxygen atoms.

13. (original): A method of stabilizing a material which is subject to degradation by actinic  
radiation by incorporating said material with the compound of claim 1.

14. (currently amended): The method of claim 13, wherein the amount of the said  
compound of claim 1 is from about 0.01 to about 20 % by weight based on the weight of the  
material to be stabilized.

15. (currently amended): The method of claim 13, wherein the material to be stabilized is  
selected from the group consisting of polyolefins, polyesters, polyethers, polyketones,  
polyamides, natural and synthetic rubbers, polyurethanes, polystyrenes, high-impact  
polystyrenes, polyacrylates, polymethacrylates, polyacetals, polyacrylonitriles,  
polybutadienes, polystyrenes, ABS acrylonitrile-butadiene-styrene, SAN (styrene  
acrylonitrile), ASA (acrylate styrene acrylonitrile), cellulosic acetate butyrate, cellulosic  
polymers, polyimides, polyamideimides, polyetherimides, polyphenylsulfide,  
PPO polyphenylene oxide, polysulfones, polyethersulfones, polyvinylchlorides,  
polycarbonates, polyketones, aliphatic polyketones, thermoplastic TPU's thermoplastic  
olefin, aminoresin cross-linked polyacrylates and polyesters, or polyisocyanate cross-linked

polyesters and polyacrylates, phenol/formaldehyde, urea/formaldehyde and melamine/formaldehyde resins, drying and non-drying alkyd resins, alkyd resins, polyester resins, acrylate resins cross-linked with melamine resins, urea resins, isocyanates, isocyanurates, carbamates, and epoxy resins, cross-linked epoxy resins derived from aliphatic, cycloaliphatic, heterocyclic and aromatic glycidyl compounds, which are cross-linked with anhydrides or amines, polysiloxanes, Michael addition polymers, amines, blocked amines with activated unsaturated and methylene compounds, ketimines with activated unsaturated and methylene compounds, polyketimines in combination with unsaturated acrylic polyacetoacetate resins, polyketimines in combination with unsaturated acrylic resins, radiation curable compositions, epoxymelamine resins, organic dyes, cosmetic products, cellulose-based paper formulations, photographic film paper, ink, fibers and combinations thereof.

16. (original): The method of claim 15, wherein the material is a polyolefin, polyamide, polyurethane, polyester or a polycarbonate.

17. (original): The method of claim 13 further comprising incorporation of one or more hindered amine light stabilizers.

18. (original): The method according to claim 17, wherein said hindered amine comprises at least one member of the group consisting of: bis(2,2,6,6-tetramethylpiperidin-4-yl) sebacate; bis(2,2,6,6-tetramethylpiperidin-4-yl)succinate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidin-4-yl)sebacate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl) n-butyl 3,5-di-tert-butyl-4-hydroxybenzylmalonate; the condensate of 1-(2-hydroxyethyl)-2,2,6,6-tetramethyl-4-hydroxypiperidine and succinic acid; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-tert-octylamino-2,6-dichloro-1,3,5-triazine; tris(2,2,6,6-tetramethylpiperidin-4-yl) nitrilotriacetate; tetrakis(2,2,6,6-tetramethylpiperidin-4-yl)- 1,2,3,4-butanetetracarboxylate; 1,1'-(1,2-ethanediyl)bis(3,3,5,5-tetramethylpiperazinone); 4-benzoyl-2,2,6,6-tetramethylpiperidine; 4-stearyloxy-2,2,6,6-tetramethylpiperidine; bis(1,2,2,6,6-pentamethylpiperidyl)-2-n-butyl-2-(2-hydroxy-3,5-di-tert-butylbenzyl)malonate; 3-n-octyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decan-2,4-dione; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)succinate; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-morpholino-2,6-dichloro-1,3,5-triazine; the condensate of 2-chloro-4,6-bis(4-n-butylamino-2,2,6,6-tetramethylpiperidyl)-

1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane; the condensate of 2-chloro-4,6-bis(4-n-butylamino-1,2,2,6,6-pentamethylpiperidyl)-1,3,5-triazine and 1,2-bis-(3-aminopropylamino)ethane; 8-acetyl-3-dodecyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decane-2,4-dione; 3-dodecyl-1-(2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1-ethanoyl-2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyrrolidine-2,5-dione; a mixture of 4-hexadecyloxy- and 4-stearyloxy-2,2,6,6-tetramethylpiperidine; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-cyclohexylamino-2,6-dichloro-1,3,5-triazine; the condensate of 1,2-bis(3-aminopropylamino)ethane, 2,4,6-trichloro-1,3,5-triazine and 4-butylamino-2,2,6,6-tetramethylpiperidine; 2-undecyl-7,7,9,9-tetramethyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane; oxo-piperanzinyl-triazines; and the reaction product of 7,7,9,9-tetramethyl-2-cycloundecyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane and epichlorohydrin.

19. (original): The method according to claim 13 further comprising incorporation of one or more additional UV absorbers wherein the additional UV absorber is a benzotriazole derivative, a triazine derivative, a benzophenone derivative, or a combination thereof.

20. (original): A method of stabilizing a material which is subject to degradation by actinic radiation by incorporating said material with the compound of claims 9, 10, 11 or 12.

21. (original): A composition comprising

(a) the compound of claim 1; and

(b) at least one other additive selected from group consisting of: UV stabilizers and antioxidants.

22. (original): The composition of claim 21 wherein said at least one other additive is selected from the group consisting of 2-(2'-hydroxyphenyl)benzotriazoles, oxamides, 2-(2-hydroxyphenyl)-1,3,5-triazines, 2-hydroxybenzophenones, sterically hindered amines and hindered phenol antioxidants.


23. (original): The composition of claim 22 wherein said at least other additive is selected from the group consisting of: 2-(2'-hydroxy-5'-methylphenyl)-benzotriazole; 2-(3',5'-di-tert-butyl-2'-hydroxyphenyl)benzotriazole; 2-(5'-tert-butyl-2'-hydroxyphenyl)benzotriazole; 2-(2'-hydroxy-5'-(1,1,3,3-tetramethylbutyl)phenyl)benzotriazole; 2-(3',5'-di-tert-butyl-2'-

hydroxyphenyl)-5-chlorobenzotriazole; 2-(3'-tert-butyl-2'-hydroxy-5'-methylphenyl)-5-chlorobenzotriazole; 2-(3'-sec-butyl-5'-tert-butyl-2'-hydroxyphenyl)-benzotriazole; 2-(2'-hydroxy-4'-octoxyphenyl)benzotriazole; 2-(3',5'-di-tert-amyl-2'-hydroxyphenyl)benzotriazole; 2-(3',5'-bis( $\alpha,\alpha$ -dimethylbenzyl)-2'-hydroxyphenyl)-benzotriazole; a mixture of 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)-carbonyl]ethyl)-2'-hydroxyphenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)carbonyl]ethyl)-2'-hydroxyphenyl)benzotriazole, 2-(3'-dodecyl-2'-hydroxy-5'-methylphenyl)benzotriazole and 2-(3'-tert-butyl-2'-hydroxy-5'-(2-isooctyloxycarbonyl)ethyl)phenylbenzotriazole; 2,2-methylenebis[4-(1,1,3,3-tetramethylbutyl)-6-benzotriazol-2-ylphenol], the transesterification product of 2-[3'-tert-butyl-5'-(2-methoxycarbonyl)ethyl)-2'-hydroxyphenyl]benzotriazole with polyethylene glycol 300; [R—CH<sub>2</sub>CH—COO(CH<sub>2</sub>)<sub>3</sub>]<sub>2</sub> B where R = 3'-tert-butyl-4'-hydroxy-5'-2H-benzotriazol-2-ylphenyl; bis(2,2,6,6-tetramethylpiperidin-4-yl) sebacate; bis(2,2,6,6-tetramethylpiperidin-4-yl)succinate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidin-4-yl)sebacate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl) n-butyl 3,5-di-tert-butyl-4-hydroxybenzylmalonate; the condensate of 1-(2-hydroxyethyl)-2,2,6,6-tetramethyl-4-hydroxypiperidine and succinic acid; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-tert-octylamino-2,6-dichloro-1,3,5-triazine; tris(2,2,6,6-tetramethylpiperidin-4-yl) nitrilotriacetate; tetrakis(2,2,6,6-tetramethylpiperidin-4-yl)-1,2,3,4-butanetetracarboxylate; 1,1'-(1,2-ethanediyl)bis(3,3,5,5-tetramethylpiperazinone); 4-benzoyl-2,2,6,6-tetramethylpiperidine; 4-stearyloxy-2,2,6,6-tetramethylpiperidine; bis(1,2,2,6,6-pentamethylpiperidyl)-2-n-butyl-2-(2-hydroxy-3,5-di-tert-butylbenzyl)malonate; 3-n-octyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decan-2,4-dione; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)succinate; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-morpholino-2,6-dichloro-1,3,5-triazine; the condensate of 2-chloro-4,6-bis(4-n-butylamino-2,2,6,6-tetramethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane; the condensate of 2-chloro-4,6-bis(4-n-butylamino-1,2,2,6,6-pentamethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane; 8-acetyl-3-dodecyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decane-2,4-dione; 3-dodecyl-1-(2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1-ethanoyl-2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyrrolidine-2,5-dione; a mixture of 4-hexadecyloxy- and 4-stearyloxy-2,2,6,6-tetramethylpiperidine; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-




yl)hexamethylenediamine and 4-cyclohexylamino-2,6-dichloro-1,3,5-triazine; the condensate of 1,2-bis(3-aminopropylamino)ethane, 2,4,6-trichloro-1,3,5-triazine and 4-butylamino-2,2,6,6-tetramethylpiperidine; 2-undecyl-7,7,9,9-tetramethyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane; oxo-piperanzinyl-triazines and the reaction product of 7,7,9,9-tetramethyl-2-cycloundecyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane and epichlorohydrin; 2,4,6-tris(2-hydroxy-4-octyloxyphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-n-octyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-(mixed iso-octyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2,4-dihydroxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2,4-bis(2-hydroxy-4-propyloxyphenyl)-6-(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-octyloxyphenyl)-4,6-bis(4-methylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-dodecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-tridecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-butyloxypropyloxy)phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-octyloxypropyloxy)-phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[4-dodecyloxy/tridecyloxy-2-hydroxypropoxy)-2-hydroxyphenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-dodecyloxypropoxy)phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-hexyloxy)phenyl-4,6-diphenyl-1,3,5-triazine; 2-(2-hydroxy-4-methoxyphenyl)-4,6-diphenyl-1,3,5-triazine; 2,4,6-tris[2-hydroxy-4-(3-butoxy-2-hydroxypropoxy)phenyl]-1,3,5-triazine; 2-(2-hydroxyphenyl)-4-(4-methoxyphenyl)-6-phenyl-1,3,5-triazine, 2,4-dihydroxybenzophenone; 2-hydroxy-4-methoxybenzophenone; 2-hydroxy-4-octyloxybenzophenone; 2-hydroxy-4-decyloxybenzophenone; 2-hydroxy-4-dodecyloxybenzophenone; 2-hydroxy-4-benzyloxybenzophenone, 4,2',4'-trishydroxybenzophenone; 2'-hydroxy-4,4'-dimethoxybenzophenone; 1,3,5-tris(2,6-dimethyl-4-tert-butyl-3hydroxybenzyl)isocyanurate; 1,3,5-tris(3,5-di-tert-butyl-4-hydroxybenzyl)isocyanurate; 1,3,5-tris(3,5-di-tert-butyl-4-hydroxybenzyl)-2,4,6-trimethylbenzene; 2,6-di-tert-butyl-4-methylphenol; 2,2'-ethylidene-bis(4,6-di-tert-butylphenol); 1,1,3-tris(5-tert-butyl-4-hydroxy-2-methylphenyl)butane; esters of  $\beta$ -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid with mono- or polyhydric alcohols; esters of  $\beta$ -(5-tert-butyl-4-hydroxy-3-methylphenyl)propionic acid with mono- or polyhydric alcohols; dimethyl-2,5-di-tert-butyl-4-hydroxybenzylphosphonate; diethyl-3,5-di-tert-butyl-4-hydroxybenzylphosphonate; dioctadecyl-3,5-di-tert-butyl-4-hydroxybenzylphosphonate; dioctadecyl-5-tert-butyl-4-hydroxy-3-methylbenzylphosphonate; and the calcium salt of the monoethyl ester of 3,5-di-tert-butyl-4-hydroxybenzylphosphonic acid; amides of  $\beta$ -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid such as N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hexamethylenediamine; N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)trimethylenediamine; and N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hydrazine.

24. (currently amended): The composition of claim 23 wherein TZ is



and T' is oxygen, ~~Y is L~~ and R<sub>1</sub> to R<sub>7</sub> and X are hydrogen.

25. (currently amended): The composition of claim 23 wherein TZ is:



and T' is oxygen, ~~Y is L~~ and R<sub>1</sub> to R<sub>7</sub> and X are hydrogen.

26. (currently amended): The composition of claim 21 further comprising a material to be stabilized selected from the group consisting of: polyolefins, polyesters, polyethers, polyketones, polyamides, natural and synthetic rubbers, polyurethanes, polystyrenes, high-impact polystyrenes, polyacrylates, polymethacrylates, polyacetals, polyacrylonitriles, polybutadienes, polystyrenes, ABS acrylonitrile-butadiene-styrene, styrene acrylonitrile, acrylate styrene acrylonitrile, cellulosic acetate butyrate, cellulosic polymers, polyimides, polyamideimides, polyetherimides, polyphenylsulfides, polyphenylene oxide, polysulfones, polyethersulfones, polyvinylchlorides, polycarbonates, polyketones, aliphatic polyketones, thermoplastic olefin-TPO's, aminoresin cross-linked polyacrylates and polyesters, polyisocyanate cross-linked polyesters and polyacrylates, phenol/formaldehyde, urea/formaldehyde and melamine/formaldehyde resins, drying and non-drying alkyd resins, alkyd resins, polyester resins, acrylate resins cross-linked with melamine resins, urea resins, isocyanates, isocyanurates, carbamates, epoxy resins, cross-linked epoxy resins derived from aliphatic, cycloaliphatic, heterocyclic and aromatic glycidyl compounds, which are cross-linked with anhydrides or amines, polysiloxanes, Michael addition polymers, amines, blocked amines with activated unsaturated and methylene compounds, ketimines with



activated unsaturated and methylene compounds, polyketimines in combination with unsaturated acrylic polyacetoacetate resins, polyketimines in combination with unsaturated acrylic resins, radiation curable compositions, epoxymelamine resins, organic dyes, cosmetic products, cellulose-based paper formulations, photographic film paper, ink, and mixtures thereof.

27. (currently amended): The composition of claim 21 wherein the amount of said compound of claim 4 to said at least one other additive if is from about 500:1 to about 1:500 by weight.

28. (original): A composition comprising

- (a) the compound of claims 9, 10, 11 or 12; and
- (b) at least one other additive selected from group consisting of: 2-(2'-hydroxy-5'-methylphenyl)-benzotriazole; 2-(3',5'-di-tert-butyl-2'-hydroxyphenyl)benzotriazole; 2-(5'-tert-butyl-2'-hydroxyphenyl)benzotriazole; 2-(2'-hydroxy-5'-(1,1,3,3-tetramethylbutyl)phenyl)benzotriazole; 2-(3',5'-di-tert-butyl-2'-hydroxyphenyl)-5-chlorobenzotriazole; 2-(3'-tert-butyl-2'-hydroxy-5'-methylphenyl)-5-chloro-benzotriazole; 2-(3'-sec-butyl-5'-tert-butyl-2'-hydroxyphenyl)-benzotriazole; 2-(2'-hydroxy-4'-octoxyphenyl)benzotriazole; 2-(3',5'-di-tert-amyl-2'-hydroxyphenyl)benzotriazole; 2-(3',5'-bis( $\alpha,\alpha$ -dimethylbenzyl)-2'-hydroxyphenyl)-benzotriazole; a mixture of 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)-carbonyl]ethyl)-2'-hydroxyphenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)-5-chloro-benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-methoxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-2'-hydroxy-5'-(2-octyloxycarbonyl)ethyl)phenyl)benzotriazole, 2-(3'-tert-butyl-5'-[2-(2-ethylhexyloxy)carbonyl]ethyl)-2'-hydroxyphenyl)benzotriazole, 2-(3'-dodecyl-2'-hydroxy-5'-methylphenyl)benzotriazole and 2-(3'-tert-butyl-2'-hydroxy-5'-(2-isooctyloxycarbonyl)ethyl)phenyl)benzotriazole; 2,2-methylenebis[4-(1,1,3,3-tetramethylbutyl)-6-benzotriazol-2-yl]phenol], the transesterification product of 2-[3'-tert-butyl-5'-(2-methoxycarbonyl)ethyl)-2'-hydroxyphenyl]benzotriazole with polyethylene glycol 300;  $[R-CH_2CH-COO(CH_2)_3]_2$  B where R = 3'-tert-butyl-4'-hydroxy-5'-2H-benzotriazol-2-ylphenyl; bis(2,2,6,6-tetramethylpiperidin-4-yl) sebacate; bis(2,2,6,6-tetramethylpiperidin-4-yl)succinate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidin-4-yl)sebacate; bis(1,2,2,6,6-pentamethylpiperidin-4-yl) n-butyl 3,5-di-tert-butyl-4-hydroxybenzylmalonate; the condensate of 1-(2-hydroxyethyl)-2,2,6,6-tetramethyl-4-hydroxypiperidine and succinic acid; the condensate of N,N'-bis(2,2,6,6-

tetramethylpiperidin-4-yl)hexamethylenediamine and 4-tert-octylamino-2,6-dichloro-1,3,5-triazine; tris(2,2,6,6-tetramethylpiperidin-4-yl) nitrilotriacetate; tetrakis(2,2,6,6-tetramethylpiperidin-4-yl)-1,2,3,4-butanetetracarboxylate; 1,1'-(1,2-ethanediyl)bis(3,3,5,5-tetramethylpiperazinone); 4-benzoyl-2,2,6,6-tetramethylpiperidine; 4-stearyloxy-2,2,6,6-tetramethylpiperidine; bis(1,2,2,6,6-pentamethylpiperidyl)-2-n-butyl-2-(2-hydroxy-3,5-di-tert-butylbenzyl)malonate; 3-n-octyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decan-2,4-dione; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)sebacate; bis(1-octyloxy-2,2,6,6-tetramethylpiperidyl)succinate; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-morpholino-2,6-dichloro-1,3,5-triazine; the condensate of 2-chloro-4,6-bis(4-n-butylamino-2,2,6,6-tetramethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane; the condensate of 2-chloro-4,6-bis(4-n-butylamino-1,2,2,6,6-pentamethylpiperidyl)-1,3,5-triazine and 1,2-bis(3-aminopropylamino)ethane; 8-acetyl-3-dodecyl-7,7,9,9-tetramethyl-1,3,8-triazaspiro[4.5]decane-2,4-dione; 3-dodecyl-1-(2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1-ethanoyl-2,2,6,6-tetramethylpiperidin-4-yl)pyrrolidin-2,5-dione; 3-dodecyl-1-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyrrolidine-2,5-dione; a mixture of 4-hexadecyloxy- and 4-stearyloxy-2,2,6,6-tetramethylpiperidine; the condensate of N,N'-bis(2,2,6,6-tetramethylpiperidin-4-yl)hexamethylenediamine and 4-cyclohexylamino-2,6-dichloro-1,3,5-triazine; the condensate of 1,2-bis(3-aminopropylamino)ethane, 2,4,6-trichloro-1,3,5-triazine and 4-butylamino-2,2,6,6-tetramethylpiperidine; 2-undecyl-7,7,9,9-tetramethyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane; oxo-piperanzinyl-triazines and the reaction product of 7,7,9,9-tetramethyl-2-cycloundecyl-1-oxa-3,8-diaza-4-oxospiro[4.5]decane and epichlorohydrin; 2,4,6-tris(2-hydroxy-4-octyloxyphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-n-octyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-(mixed iso-octyloxyphenyl))-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2,4-dihydroxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2,4-dihydroxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-propyloxyphenyl)-6-(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-octyloxyphenyl)-4,6-bis(4-methylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-dodecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-tridecyloxyphenyl)-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-butyloxypropyloxy)phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-octyloxypropyloxy)-phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[4-dodecyloxy/tridecyloxy-2-hydroxypropoxy]-2-hydroxyphenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-[2-hydroxy-4-(2-hydroxy-3-dodecyloxypropoxy)phenyl]-4,6-bis(2,4-dimethylphenyl)-1,3,5-triazine; 2-(2-hydroxy-4-hexyloxy)phenyl-4,6-diphenyl-1,3,5-triazine; 2-(2-hydroxy-4-methoxyphenyl)-4,6-diphenyl-1,3,5-triazine; 2,4,6-tris[2-hydroxy-4-(3-butoxy-2-hydroxypropoxy)phenyl]-1,3,5-triazine; 2-(2-hydroxyphenyl)-4-(4-methoxyphenyl)-6-phenyl-1,3,5-triazine, 2,4-dihydroxybenzophenone; 2-hydroxy-4-methoxybenzophenone; 2-


hydroxy-4-octyloxybenzophenone; 2-hydroxy-4-decyloxybenzophenone; 2-hydroxy-4-dodecyloxybenzophenone; 2-hydroxy-4-benzyloxybenzophenone, 4,2',4-trishydroxybenzophenone; 2'-hydroxy-4,4'-dimethoxybenzophenone; 1,3,5-tris( 2,6-dimethyl-4-tert-butyl-3hydroxybenzyl)isocyanurate; 1,3,5-tris(3,5-di-tert-butyl-4-hydroxybenzyl)isocyanurate; 1,3,5- tris(3,5- di-tert-butyl-4-hydroxybenzyl) -2,4,6-trimethylbenzene; 2,6-di-tert-butyl-4-methylphenol; 2,2'-ethylidene-bis(4,6-di-tert-butylphenol); 1,1,3-tris(5-tert-butyl-4-hydroxy-2-methylphenyl)butane; esters of  $\beta$ -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid with mono- or polyhydric alcohols; esters of  $\beta$ -(5-tert-butyl-4-hydroxy-3-methylphenyl)propionic acid with mono- or polyhydric alcohols; dimethyl-2,5-di-tert-butyl-4-hydroxybenzylphosphonate; diethyl-3,5-di-tert-butyl-4-hydroxybenzylphosphonate; dioctadecyl-3,5-di-tert-butyl-4-hydroxybenzylphosphonate; dioctadecyl-5-tert-butyl-4-hydroxy-3-methylbenzylphosphonate; and the calcium salt of the monoethyl ester of 3,5-di-tert-butyl-4-hydroxybenzylphosphonic acid; amides of  $\beta$ -(3,5-di-tert-butyl-4-hydroxyphenyl)propionic acid such as N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hexamethylenediamine; N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)trimethylenediamine; and N,N'-bis(3,5-di-tert-butyl-4-hydroxyphenylpropionyl)hydrazine.

29. (currently amended): The composition of claim 28 wherein the amount of said compound to said at least one other additive ~~if~~ is from about 500:1 to about 1:500 by weight.

30. (currently amended): The composition of claim 28 further comprising a material to be stabilized selected from the group consisting of polyolefins, polyesters, polyethers, polyketones, polyamides, natural and synthetic rubbers, polyurethanes, polystyrenes, high-impact polystyrenes, polyacrylates, polymethacrylates, polyacetals, polyacrylonitriles, polybutadienes, polystyrenes, ABS acrylonitrile-butadiene-styrene, ~~SAN~~-(styrene acrylonitrile), ~~ASA~~-(acrylate styrene acrylonitrile), cellulosic acetate butyrate, cellulosic polymers, polyimides, polyamideimides, polyetherimides, polyphenylsulfide, ~~PPO~~, polyphenylene oxide, polysulfones, polyethersulfones, polyvinylchlorides, polycarbonates, polyketones, aliphatic polyketones, thermoplastic olefin~~TPU~~'s, aminoresin cross-linked polyacrylates and polyesters, or polyisocyanate cross-linked polyesters and polyacrylates, phenol/formaldehyde, urea/formaldehyde and melamine/formaldehyde resins, drying and non-drying alkyd resins, alkyd resins, polyester resins, acrylate resins cross-linked with melamine resins, urea resins, isocyanates, isocyanurates, carbamates, and epoxy resins, cross-linked epoxy resins derived from aliphatic, cycloaliphatic, heterocyclic

and aromatic glycidyl compounds, which are cross-linked with anhydrides or amines, polysiloxanes, Michael addition polymers, amines, blocked amines with activated unsaturated and methylene compounds, ketimines with activated unsaturated and methylene compounds, polyketimines in combination with unsaturated acrylic polyacetoacetate resins, polyketimines in combination with unsaturated acrylic resins, radiation curable compositions, epoxymelamine resins, organic dyes, cosmetic products, cellulose-based paper formulations, photographic film paper, ink, fibers and combinations thereof.


31. (currently amended): A compound having the Formula



wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> are the same or different and each is hydrogen, halogen, alkyl of 1 to 24 carbon atoms, haloalkyl of 1 to 24 carbon atoms, aryl of 6 to 24 carbon atoms, alkenyl of 2 to 24 carbon atoms, acyl of 1 to 24 carbon atoms, aralkyl of 7 to 24 carbon atoms, aracyl of 6 to 24 carbon atoms, OR, NRR', CONRR', OCOR, CN, SR, SO<sub>2</sub>R, SO<sub>3</sub>H, SO<sub>3</sub>M, wherein M is an alkali metal, R and R' are the same or different and each is hydrogen, alkyl of 1 to 24 carbon atoms, haloalkyl of 1 to 24 carbon atoms, aryl of 6 to 24 carbon atoms, alkenyl of 2 to 24 carbon atoms, acyl of 1 to 24 carbon atoms, cycloalkyl of 1 to 24 carbon atoms, cycloacyl of 5 to 24 carbon atoms, aralkyl of 7 to 24 carbon atoms, or aracyl of 6 to 24 carbons atoms, and optionally with either of R<sub>1</sub> and R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub>, taken together being a part of a saturated or unsaturated fused carbocyclic ring optionally containing O, N, or S atoms in the ring;

Z' is a halogen,

and W is



or a halogen.

32-34. (cancelled)